

<http://www.ncbi.nlm.nih.gov/pubmed/2195843>(**Bone Density Changes in Osteoporosis prone Women Exposed to Pulsed Electromagnetic Fields / PEMFs (1990)**). "The data suggest that properly applied PEMFs, if scaled for whole-body use, may have clinical application in the prevention and treatment of osteoporosis .")□□

Bone density changes in osteoporosis-prone women exposed to pulsed electromagnetic fields (PEMFs).

Tabrah F, Hoffmeier M, Gilbert F Jr, Batkin S, Bassett CA.

Author information

Abstract

To determine the effect of a 72 Hz pulsating electromagnetic field (PEMF) on bone density of the radii of osteoporosis-prone women, the nondominant forearms of 20 subjects were exposed to PEMF 10 h daily for a period of 12 weeks. Bone density before, during, and after the exposure period was determined by use of a Norland-Cameron bone mineral analyzer. Bone mineral densities of the treated radii measured by single-photon densitometry increased significantly in the immediate area of the field during the exposure period and decreased during the following 36 weeks. A similar but weaker response occurred in the opposite arm, suggesting a "cross-talk" effect on the nontreated radii, from either possible arm proximity during sleep or very weak general field effects. The data suggest that properly applied PEMFs, if scaled for whole-body use, may have clinical application in the prevention and treatment of osteoporosis.

PMID: 2195843 [PubMed - indexed for MEDLINE]